























































HAZARDOUS DRUGS TO WATCH FOR IN VETERINARY PRACTICE

Drug	Recommended Precautions	Who Should Take Precautions
Apomorphine	  + Wear a face mask if handling powder.	
Azathioprine	 	
Bedinvetmab Frunevetmab	 + Use PPE to minimize the risk of accidental self-injection.	
Chemotherapeutics	  + Wear gloves when handling the drug and when handling bodily fluids or litter from treated animals. Review the specific handling requirements for each drug.	
Chloramphenicol	  + Avoid crushing or cutting pills. If an animal vomits after administration, wear gloves while cleaning.	
Cyclosporine	 + Wash or rinse exposed skin or eyes.	



Drug	Recommended Precautions	Who Should Take Precautions
Dexmedetomidine oromucosal gel	 + Wear gloves when handling the syringe, administering the drug, or coming into contact with the treated dog's mouth after application.	
Diethylstilbestrol	 + Avoid crushing or cutting pills and inhaling dust. Wash or rinse exposed skin or eyes.	
	 + Any person who is pregnant, breastfeeding, or trying to conceive should avoid handling.	
Estriol	 + Avoid crushing or cutting pills and inhaling dust. Wash or rinse exposed skin or eyes.	
	 + Any person who is pregnant, breastfeeding, or trying to conceive should avoid handling.	
Finasteride		
Methimazole	 + Wear gloves when handling the bodily fluids or litter from treated animals.	
Misoprostol	 + Avoid direct contact with bodily fluids or litter of treated animals.	
Mitotane	 + Wash or rinse exposed skin or eyes. Wear gloves when handling bodily fluids, feces, or litter.	



Drug	Recommended Precautions	Who Should Take Precautions
Oclacitinib		
Ropinirole	  + Wear protective eyewear.	
Tacrolimus	  + Avoid contact with skin or mucous membranes. Wear gloves when handling bodily fluids, feces, or litter.	
Toceranib	 + Wear gloves when handling bodily fluids, feces, litter, or broken or moistened tablets. Keep medication and bodily fluids of treated animals out of reach of children.	
Transdermal fentanyl	 + Take additional steps if the patch doesn't stick to the patient's skin (eg, adhesives, bandages). Double-check patch placement regularly. Dispose of used patch in toilet.	
Trilostane	 + Do not empty capsule contents. Do not attempt to divide capsules.	
Zonisamide		



Wear gloves while handling



Wash hands after handling



Avoid handling



All team members



Any person who is pregnant, breastfeeding, or trying to conceive

*This is not an all-inclusive list. The complete list of hazardous drugs is available at [cdc.gov/niosh/docs/2016-161/pdfs/2016-161.pdf](https://www.cdc.gov/niosh/docs/2016-161/pdfs/2016-161.pdf)



Scan the QR code or visit plumbs.com to see why thousands of veterinary professionals rely on Plumb's™ for trusted prescribing support.